

Eur J Vasc Endovasc Surg 17, 373–389 (1999)  
Article No. ejvs.1998.0759

## REVIEW ARTICLE

# Cellular and Molecular Mechanisms of Inflammation and Thrombosis

P. Tan<sup>1</sup>, F. W. Lusinskas<sup>2</sup> and S. Homer-Vanniasinkam<sup>\*3</sup>

<sup>1</sup>Sheffield Vascular Institute, Northern General Hospital, Sheffield, U.K., <sup>2</sup>Vascular Research Division, Departments of Pathology, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts, U.S.A. and <sup>3</sup>Vascular Surgical Unit, The General Infirmary at Leeds, U.K.

*In the last 20 years, the cellular and molecular mechanisms of inflammation and thrombosis have been characterised. These are essentially cell adhesion processes which are regulated by vascular endothelium. Many of the cell adhesion molecules and leucocyte chemoattractants expressed and generated at sites of inflammation have been sequenced and cloned. These inflammatory molecules work together in concert to mediate the adhesion between leucocytes, platelets and vascular endothelium which occurs during the occlusive, thromboembolic, reperfusion and septic complications of atherosclerotic and diabetic vascular diseases. This review aims to summarise our current understanding of the molecular basis of these disorders and the therapeutic implications.*

**Key Words:** Cell adhesion molecules; Chemokines; Neutrophils; Inflammation; Atherosclerosis; Thrombosis.

## Introduction

The human body reacts to injurious stimuli by mounting an inflammatory response. In its acute phase, transvascular shifts of fluid, plasma proteins and leucocytes occur in the microvasculature of the injured tissue. Vascular injury induces the activation of both platelets and coagulation proteins which work together to maintain haemostasis. These physiological processes are intended to neutralise the injurious stimulus, limit tissue injury and initiate healing and repair, and this was recognised by a surgeon, John Hunter, as early as the 19th century.<sup>1</sup>

The importance of vascular endothelium as a regulatory organ in immunity and haemostasis has become well established in the last two decades.<sup>2</sup> Normally, vascular endothelium is in an “unactivated” state. However, at sites of injury and inflammation, endothelium becomes locally “activated”, either by the injurious stimulus itself, or by inflammatory cytokines generated in response to the injurious stimulus. Activated vascular endothelium expresses a number of

membrane-bound pro-inflammatory molecules, in particular, leucocyte-endothelial cell adhesion molecules (CAM). It also secretes soluble factors which attract peripheral blood leucocytes and induce coagulation, thus resulting in the leucocyte recruitment and platelet thrombosis observed at sites of inflammation.

Inflammatory cytokines are small peptides secreted primarily by activated tissue macrophages and lymphocytes in response to injurious stimuli, such as endotoxin, immune complexes, physical and chemical injury. Interleukin-1 $\beta$  (IL-1 $\beta$ ) and tumour necrosis factor  $\alpha$  (TNF  $\alpha$ ) are the main cytokines that mediate the inflammatory response.<sup>3</sup> They have a wide range of actions on different cells but their main effects are to induce the acute-phase reaction and to activate vascular endothelium, leucocytes, platelets and fibroblasts, thus, initiating the cascade of vascular, cellular and humoral events which together comprise the inflammatory response.

In this review, we will focus on the role of non-lymphoid vascular endothelium, leucocyte-endothelial CAM, chemoattractants, leucocytes and platelets as the primary molecular and cellular components of the inflammatory response. The currently understood molecular mechanisms of atherogenesis and thrombosis will also be summarised. We will conclude by discussing possible therapeutic applications.

\* Please address all correspondence to: S. Homer-Vanniasinkam, The General Infirmary at Leeds, Great George Street, Leeds LS1 3EX, U.K.

## Vascular Endothelium

Although a mere cellular monolayer, vascular endothelium contains the entire circulation and forms a vast surface area of interface between the components of the circulation (including leucocytes, platelets and plasma proteins) and all of the body's tissues. It performs a wide variety of functions including the regulation of vascular permeability and vascular tone, wound healing, immunity and haemostasis, thus, serving as a "gatekeeper" for the traffic of water and solutes, bioactive proteins and lipids, and peripheral blood leucocytes, between the circulation and interstitial tissues.

As a regulatory organ, vascular endothelium serves both sensory and effector functions, bearing receptors for a wide range of cytokines, chemotactic and vasoactive peptides and lipids, and CAM, as well as itself being able to express and elaborate many of these immunomolecules. Selective spatial and temporal expression of these molecules by vascular endothelium provides a means by which inflammatory and thrombotic processes may be regulated. Vascular endothelium continually senses chemical, physical and biomechanical alterations in its surroundings and transduces these into biological responses. For example, changes in blood flow have been shown to induce alterations in the synthesis of several vascular endothelial-dependent factors, such as prostacyclin, nitric oxide and intercellular adhesion molecule-1.<sup>4-6</sup>

### *Regulatory role in leucocyte traffic*

*In vivo* studies in leucocyte-endothelial adhesion in the microcirculation and *in vitro* studies performed under flow conditions designed to simulate microvascular blood flow indicate that unactivated vascular endothelium is hypo-adhesive for leucocytes, by virtue of minimal expression of leucocyte-endothelial CAM.<sup>7</sup> However, stimuli such as endotoxin,<sup>8</sup> inflammatory cytokines,<sup>3</sup> histamine and thrombin<sup>9</sup> induce local endothelial activation or stimulation. Activated or stimulated vascular endothelium is hyperadhesive for leucocytes due to upregulated expression of CAM such as E-selectin,<sup>10</sup> intercellular adhesion molecule-1<sup>11</sup> and P-selectin.<sup>9</sup> Additionally, activated endothelial cells express and secrete chemotactic factors such as platelet-activating factor<sup>12</sup> and interleukin-8<sup>13</sup> which may activate adherent leucocytes and direct their migration across the endothelium and within the inflamed tissues.

Multiple inflammatory response genes which are

regulated by cytoplasmic transcription factors, such as nuclear factor  $\kappa$ B,<sup>14</sup> are induced during endothelial activation. Thus, endothelial activation is a process which occurs over a period of hours being dependent on gene expression and *de novo* protein synthesis. For example, the endothelial CAM, E-selectin, which is believed to be a "silent" gene under normal circumstances, is synthesised and expressed by activated endothelial cells, and becomes detectable between 1 and 24 hours after activation.<sup>15</sup> In contrast, thrombogenic mediators, such as histamine and thrombin, induce endothelial stimulation which occurs rapidly (within minutes) and independently of gene expression and protein synthesis. P-selectin is expressed transiently during endothelial stimulation.

### *Regulatory role in haemostasis*

Unactivated endothelial cells are remarkable in being the only cells known to be actively antithrombotic. Firstly, they form an insulating barrier between thrombogenic subendothelial matrix components, and platelets and coagulation proteins in the circulation. Secondly, unactivated endothelial cells secrete prostacyclin and nitric oxide which inhibit platelet aggregation,<sup>4,5</sup> as well as protein S<sup>16</sup> and tissue plasminogen activator<sup>17</sup> which inhibit coagulation.

However, activated vascular endothelium is prothrombotic. It secretes tissue factor which triggers the coagulation cascade<sup>18,19</sup> and expresses increased levels of plasminogen activator inhibitor-1 which inhibits fibrinolysis.<sup>20</sup> Vascular injury may expose subendothelial matrix components such as collagen and von Willebrand factor (vWF) which promote and support platelet adhesion and aggregation.

Thus, at rest, vascular endothelium maintains blood in its fluid form but, in response to vascular injury, it becomes locally prothrombotic to seal off rapidly any breach in the integrity of the vasculature.

## Leucocyte-endothelial Cell Adhesion Molecules

The histological hallmark of acute inflammation is the infiltration of interstitial tissues by peripheral blood leucocytes that consists in the early phases predominantly of neutrophils and, later on, mononuclear leucocytes. The ability of peripheral blood leucocytes to extravasate at sites of inflammation was described over a century ago,<sup>21</sup> but only in the past decade have the molecules responsible for leucocyte adhesion and

transendothelial migration (also called diapedesis, emigration) been identified and cloned, and their mechanisms of action defined.<sup>22–24</sup>

Leucocyte-endothelial cell adhesion molecules (CAM) are proteins expressed on the surface of leucocytes and platelets and the surface of vascular endothelium. They mediate the adhesive interactions between circulating leucocytes and endothelium leading to leucocyte recruitment into tissues, and between platelets and endothelium resulting in thrombosis. The role of CAM in leucocyte trafficking in health and disease has been established by *in vitro* experiments with cultured vascular endothelium and isolated leucocytes<sup>25–27</sup> and by *in vivo* experiments using animal models.<sup>28,29</sup> The generation of function-blocking monoclonal antibodies against individual CAM has enabled the various steps of leucocyte-endothelial adhesion to be characterised. More recently, experiments using genetically-engineered mutant mice lacking one or more CAM ("knockout mice") have further clarified their respective roles in leucocyte trafficking and immunity.<sup>30–33</sup> Furthermore, the importance of CAM in host defence is vividly illustrated by two rare autosomal recessively inherited diseases, leucocyte adhesion deficiency (LAD) Types I & II.<sup>34,35</sup> LAD I patients have deficient neutrophil diapedesis and phagocytosis due to molecular defects in their leucocyte integrins. Similarly, LAD II patients exhibit deficient neutrophil adhesion due to a metabolic defect in the synthesis of selectin ligands. LAD patients are markedly immunodeficient and suffer from severe, recurrent and often fatal bacterial infections.

To date, endothelial-leucocyte CAM have been classified according to structural homology into 4 families: the selectins, integrins, mucin-like CAM and the immunoglobulin superfamily (Table 1). The salient characteristics only of these CAM will be discussed below as more detailed reviews already exist.<sup>22–24,36</sup>

### Selectins

Selectins are transmembrane glycoproteins expressed by leucocytes, and activated endothelium and platelets. Their primary function appears to be to mediate the initial attachment and rolling (discussed below) adhesive interactions that occur under pathophysiological conditions of blood flow between leucocytes and postcapillary venular endothelium at sites of inflammation.<sup>24</sup> There are three members of the selectin family.

CD62L or L-selectin ("L" for leucocyte) is constitutively expressed on the surface of most leuco-

cytes.<sup>37</sup> Neutrophils and monocytes bear high surface levels of L-selectin that is lost or shed by proteolytic cleavage on contact with activated endothelium or chemoattractants.<sup>27,38</sup> A basal level of shedding appears to result in high levels of functionally active L-selectin in plasma<sup>39</sup> which may reach levels sufficient to partially inhibit leucocyte-endothelial adhesion.<sup>40</sup> However, *in vitro*, L-selectin shedding *per se* neither enhances nor inhibits neutrophil-endothelial adhesion and transmigration.<sup>41</sup>

CD62P or P-selectin ("P" for platelet) is normally not expressed but sequestered in intracellular storage granules in vascular endothelial cells (in Weibel-Palade bodies) and platelets (in  $\alpha$ -granules).<sup>9</sup> Thrombogenic and inflammatory mediators such as histamine, thrombin, and leucotrienes C & D induce rapid translocation of P-selectin to the surface of platelets and vascular endothelial cells. *In vitro*, the expression of P-selectin is rapid and short lived (expressed then downregulated within minutes).

CD62E or E-selectin ("E" for endothelial) is expressed exclusively by activated vascular endothelial cells.<sup>15</sup> Induction of the E-selectin gene in cultured human umbilical vein endothelial cells by inflammatory stimuli, such as endotoxin, IL-1 $\beta$  or TNF- $\alpha$ , induces gene transcription and surface expression of the E-selectin protein which is detectable within an hour of endothelial activation. *In vitro*, peak E-selectin expression occurs around 4–6 h and becomes undetectable after 12–24 h. However, more prolonged expression of E-selectin may exist *in vivo* in chronic inflammatory diseases.<sup>42,43</sup>

The physiological counter-receptors of selectins have not been fully characterised, but they appear to be carbohydrate structures presented by glycoproteins on the surface of vascular endothelium, leucocytes and platelets.<sup>24,44</sup>

### Mucin-like proteins

P-selectin glycoprotein ligand-1 (PSGL-1) is the best characterised selectin counter-receptor.<sup>45,46</sup> PSGL-1 is present on neutrophils, monocytes and lymphocytes and binds L-selectin, E-selectin and P-selectin.<sup>47–49</sup> Neutrophil-neutrophil adhesive interactions that occur during neutrophil aggregation as well as during initial attachment and rolling (discussed below) of neutrophils on activated vascular endothelium are mediated by PSGL-1–L-selectin bonds.<sup>47,50</sup> Current evidence suggests that another ligand for L-selectin may be expressed by activated non-lymphoid vascular

**Table 1. Endothelial-leucocyte cell adhesion molecules.**

Family	Members	Cluster designation	Cellular distribution	Counter-receptor/ligand
Selectins	E-selectin	CD62E	Endothelium	PSGL-1, ?other carbohydrate-bearing structure(s) on leucocytes
	L-selectin	CD62L	Leucocytes	PSGL-1, ?inducible carbohydrate-bearing structure(s) on endothelium
	P-selectin	CD62P	Endothelium, platelets	PSGL-1, ?other carbohydrate-bearing structure(s) on leucocytes
Mucin-like	PSGL-1	CD162	All blood leucocytes	E-, L- and P-selectin
$\beta_1$ integrins	$\alpha_4\beta_1$	CD29	Monocytes, lymphocytes	VCAM-1
$\beta_2$ integrins	LFA-1	CD11a/CD18	Leucocytes	ICAM-1, ICAM-2, ICAM-3
	Mac-1	CD11b/CD18	Monocytes, neutrophils	ICAM-1, fibrinogen
$\beta_7$ integrins	p150,95	CD11c/CD18	Monocytes, neutrophils	?
	$\alpha_4\beta_7$		Lymphocytes	VCAM-1
Ig	ICAM-1	CD54	Endothelium, leucocytes, epithelial cells, fibroblasts, other cell lines	LFA-1, Mac-1
	ICAM-2	CD102	Endothelium	LFA-1
	ICAM-3	CD50	Leucocytes	LFA-1
	VCAM-1	CD106	Endothelium, smooth muscle cells	$\alpha_4\beta_1$ , $\alpha_4\beta_7$
	PECAM-1	CD31	Endothelium, leucocytes, platelets	PECAM-1

endothelium, although it has yet to be identified.<sup>24,51</sup> Other mucin-like proteins such as GlyCAM-1, CD34 and ESL-1, which bind selectins, have been described in animal models, but except for CD34 these have not been shown to be counter-receptors for selectins in humans.<sup>24</sup>

### *Leucocyte integrins*

Integrins are transmembrane glycoproteins widely expressed on many cells.<sup>34,52</sup> They are heterodimers of an  $\alpha$  subunit non-covalently linked to a  $\beta$  subunit and have been classified into subfamilies according to their  $\beta$  chain (8  $\beta$  chains have been identified to date).

The main integrin subfamilies relevant to the role of leucocytes and platelets in inflammation and thrombosis are  $\beta_1$ ,  $\beta_2$ ,  $\beta_3$  and  $\beta_7$ .  $\beta_1$  integrins are expressed by a wide variety of vascular and non-vascular cells, the  $\beta_1$  chain (CD29) combining with a variety of  $\alpha$  chains.  $\beta_3$  integrins or cytoadhesins are expressed by platelets, endothelial cells and smooth muscle cells (the  $\beta_3$  chain being designated CD61).  $\beta_1$  and  $\beta_3$  integrins will be briefly discussed later in the context of platelets and thrombosis. However, the integrin subfamily most relevant to endothelial-leucocyte adhesion is the  $\beta_2$  (CD18) or leucocyte integrin subfamily.

$\beta_2$  integrins are expressed exclusively by leucocytes. The  $\beta_2$  integrin subfamily has three members all of

which have CD18 as their common  $\beta$  chain: lymphocyte function associated antigen-1 (LFA-1) or  $\alpha_L\beta_2$  (CD11a/CD18), Mac-1 or  $\alpha_M\beta_2$  (CD11b/CD18), and p150,95 or  $\alpha_X\beta_2$  (CD11c/CD18). To date, neutrophils are known to express only  $\beta_2$  integrins while monocytes and lymphocytes express both  $\beta_1$  and  $\beta_2$  integrins.<sup>36</sup>

Unactivated neutrophils express high surface levels of LFA-1 and Mac-1 but minimal amounts of p150,95.<sup>53</sup> Mac-1 is also stored within intracellular peroxidase-negative granules.<sup>54</sup> On activation by inflammatory mediators such as chemoattractants, *in vitro*, the surface expression of Mac-1 is upregulated several-fold by translocation from cytoplasmic granules while that of LFA-1 remains relatively unchanged.<sup>55-57</sup> However, these newly-expressed Mac-1 molecules may not all be functionally competent, as further conformational change with the expression of activation neo-epitopes appears to be necessary for ligand binding. Expression of activation neo-epitopes which confer the ability to bind ligand has been described for both LFA-1 and Mac-1.<sup>55,58</sup>

LFA-1 counter-receptors include intercellular adhesion molecule-1 (ICAM-1), ICAM-2 and ICAM-3<sup>59-61</sup> while Mac-1 binds to ICAM-1, fibrinogen<sup>55,62</sup> and opsonic complement fragment (iC2b).<sup>63</sup> Mac-1 is also believed to mediate the binding of neutrophils to plastic and glass, and aggregation with other neutrophils.<sup>53</sup> The counter-receptor(s) of p150,95 are less certain, but it may be a complement receptor.<sup>64</sup>



*Immunoglobulin superfamily*

CAM belonging to the immunoglobulin (Ig) superfamily consist of cell surface proteins which are related structurally by possessing varying numbers of Ig-like domains. There are five members of the Ig-like family which mediate endothelial-leucocyte adhesion: intercellular adhesion molecule-1 (ICAM-1), ICAM-2, ICAM-3, vascular cell adhesion molecule-1 (VCAM-1) and platelet-endothelial cell adhesion molecule-1 (PECAM-1).

ICAM-1 (CD54) is widely expressed by many cell types including vascular endothelium, fibroblasts, epithelial cells, skeletal myocytes, and some leucocytes but not neutrophils.<sup>65</sup> It is constitutively expressed at low levels by vascular endothelium but its expression is upregulated several-fold by endotoxin and inflammatory cytokines.<sup>11</sup> Its counter-receptors include LFA-1, Mac-1, rhinovirus<sup>66</sup> and *Plasmodium falciparum*.<sup>67</sup> Soluble ICAM-1 detected in human plasma<sup>39</sup> has been proposed as a means by which the body modulates ICAM-1 mediated adhesion (by binding to its counter-receptors such as LFA-1 and Mac-1 on leucocytes). However, the concentrations of soluble ICAM-1 necessary to competitively inhibit ICAM-1–LFA-1 adhesion exceed those normally found in plasma and therefore, it is unlikely that soluble ICAM-1 would antagonise ICAM-1 mediated adhesion *in vivo*.<sup>68</sup>

ICAM-2 (CD102) is constitutively expressed by vascular endothelium and some mononuclear leucocytes. Its level of expression by vascular endothelium is not cytokine-inducible.<sup>69</sup> ICAM-3 (CD50), on the other hand, is highly expressed on leucocytes but not vascular endothelium.<sup>61</sup> Both ICAM-2 and ICAM-3 have been shown to bind LFA-1 but their role in leucocyte-endothelial adhesion remains uncertain.<sup>60,61,69</sup>

VCAM-1 (CD106) is expressed mainly by vascular endothelium but may also be expressed by vascular smooth muscle cells, dendritic cells and synovial cells.<sup>70–72</sup> VCAM-1 expression is focally upregulated by inflammatory and atherogenic stimuli such as IL-1 $\beta$ , interleukin-4 and lysophosphatidylcholine (a component of atherogenic lipoproteins).<sup>73–75</sup> Counter-receptors identified for VCAM-1 to date include the  $\beta_1$  and  $\beta_7$  integrins, VLA-4 ( $\alpha_4\beta_1$ ) and  $\alpha_4\beta_7$ .<sup>76,77</sup> Neutrophils do not express  $\beta_1$  and  $\beta_7$  integrins and cannot bind to VCAM-1.

PECAM-1 (CD31) is constitutively expressed by vascular endothelium, neutrophils, monocytes, some lymphocytes and platelets.<sup>23</sup> Its expression on vascular endothelium is characteristic, being concentrated at intercellular junctions<sup>78</sup> where it mediates adhesion (CD31–CD31) between vascular endothelial cells.

CD31 expression is not upregulated by inflammatory cytokines, although its distribution on the surface of endothelial cells may be altered by interferon  $\gamma$  and TNF $\alpha$ .<sup>79,80</sup> It also mediates adhesion between myeloid cells (neutrophils and monocytes) and platelets as well as between myeloid cells and vascular endothelium. It appears to be important in mediating neutrophil and monocyte transmigration between intercellular junctions of normal and activated vascular endothelium.<sup>81,82</sup> Recently, CD31 has been shown to undergo adhesion to the integrin  $\alpha_v\beta_3$ .<sup>83</sup>

*Platelet glycoproteins*

Platelets are crucial to the body's normal inflammatory response because of their ability to form haemostatic platelet plugs very rapidly at sites of vascular injury. They also play an important role in initiating healing processes, by virtue of the growth and chemotactic factors stored in their alpha granules and released on activation. The surfaces of these tiny non-nucleated cells are densely decorated with  $\beta_1$  and  $\beta_3$  integrins and other platelet glycoproteins (gp) which mediate the two platelet reactions that are crucial for normal haemostasis, namely, platelet adhesion and platelet aggregation. Platelets adhere to subendothelial matrix proteins exposed by vascular injury (platelet adhesion). Then, on activation, by the process of platelet adhesion and/or by agonists such as adenine diphosphate (ADP), adrenaline and thrombin, more platelets are recruited and begin to adhere to each other (platelet aggregation) and to form a haemostatic plug onto which fibrin may be deposited. Leucocyte recruitment mediated by P-selectin appears to be involved in this process of fibrin deposition.<sup>84</sup>

Platelet adhesion is mediated by  $\alpha_2\beta_1$  (binds collagen),  $\alpha_5\beta_1$  (binds fibronectin),  $\alpha_6\beta_1$  (binds laminin) and  $\alpha_{IIb}\beta_3$  (also called, gpIIb/IIIa) as well as other glycoproteins, such as gpIb (binds vWF).<sup>85,86</sup> Platelet aggregation, however, is mediated exclusively by the  $\beta_3$  integrin,  $\alpha_{IIb}\beta_3$  (gpIIb/IIIa). gpIIb/IIIa binds to a number of soluble proteins including fibrinogen and vWF but becomes functional (by "uncovering" of ligand binding sites which lie hidden on unactivated platelets) only on platelet activation.<sup>87,88</sup> Fibrinogen and vWF form crossbridges between gpIIb/IIIa molecules on adjacent platelets, thereby resulting in platelet aggregation. Blockade of gpIIb/IIIa totally inhibits platelet aggregation.

**Chemoattractants**

The process of "chemotaxis" may be defined as the unidirectional locomotion of a cell in response to a

Table 2. Leucocyte chemoattractants.

Family	Member	Source	Target cell(s)
"Classical"	C5a	Plasma protein cleavage product	Monocyte, neutrophils, basophils, eosinophils
	fMLP	Bacteria	Monocytes, neutrophils
	PAF	Leucocytes	Monocytes, neutrophils, eosinophils
	LTB <sub>4</sub>	Leucocytes	Neutrophils
C-X-C chemokines	IL-8	Most somatic cells	Neutrophils, basophils
	GRO- $\alpha$ , - $\beta$ , - $\gamma$	Most somatic cells	Neutrophils, basophils
	IP-10	Most somatic cells	Monocytes
	PF4	Platelets	Monocytes, neutrophils
C-C chemokines	MCP-1, -2, -3	Most somatic cells	Monocytes, basophils, specific lymphocyte subsets
	MIP-1 $\alpha$	Most somatic cells	Monocytes, basophils, eosinophils, specific lymphocyte subsets
	MIP-1 $\beta$	Most somatic cells	Monocytes, specific lymphocyte subsets
	RANTES	Most somatic cells	Monocytes, basophils, eosinophils, specific lymphocyte subsets

concentration gradient of a chemoattractant substance. Leukocytes, in particular, have long been observed to exhibit chemotaxis.<sup>21</sup> Chemical substances or microorganisms placed at extravascular sites, such as the anterior ocular chamber or the peritoneal cavity, induce the accumulation of large numbers of peripheral blood leucocytes at these sites as part of an inflammatory response.

To date, a wide range of chemotactic molecules have been identified which attract and activate leucocytes.<sup>89-91</sup> Furthermore, specific chemotactic receptors found on the leucocyte surface have now been cloned.<sup>92,93</sup> Presumably, by using these receptors leucocytes are able "to sense" small concentration differences of chemotactic substances across their diameter and then crawl in the direction of increasing concentration.<sup>94</sup>

#### "Classical" chemoattractants

A variety of bioactive peptides and lipids of endogenous and exogenous origin were identified in the 1970s and found to be leucocyte chemoattractants. Activated neutrophils produce an oxidised phospholipid leucotriene B<sub>4</sub> (LTB<sub>4</sub>),<sup>95</sup> as well as an ether phospholipid, platelet-activating factor (PAF),<sup>96,97</sup> both of which are chemotactic for neutrophils. PAF-like molecules are also generated by activated vascular endothelium in a membrane-bound form.<sup>98</sup> PAF, apart from being a leucocyte chemoattractant, has diverse biological activities including enhanced adhesion of platelets and leucocytes to endothelium, vasoconstriction and bronchoconstriction.<sup>96,99</sup> Both C5a anaphylatoxin, a plasma peptide generated by complement activation, and its cleavage product, C5a<sub>des arg</sub>, are potent chemoattractants for neutrophils,<sup>100</sup>

basophils, eosinophils and monocytes. fMLP, a bacterial oligopeptide, has also been shown to induce chemotaxis in neutrophils and monocytes.<sup>101</sup> It should be noted that these chemoattractants neither exhibit specificity of chemoattraction for any particular leucocyte subset nor are their biological activities confined to inducing leucocyte chemotaxis alone. PAF, as its name suggests, is a potent activator of platelets and leucocytes as well as having vasoactive properties.<sup>99</sup>

#### Chemokines

Chemokines (contraction of *chemotactic cytokines*) are a family of small pro-inflammatory peptides with potent leucocyte chemoattractant and activating properties.<sup>90,102</sup> To date, they have been classified into two main subfamilies according to amino acid sequence, their gene locus and the leucocyte subset on which they primarily exert chemotactic activity (Table 2). This specificity of chemoattraction of a particular leucocyte subset is a key feature of chemokines, in contrast to "classical" chemoattractants discussed above.

Chemokines are mostly inducible molecules, that is, produced and secreted only on cellular activation by endotoxin or inflammatory cytokines or in response to agents such as crystals and viruses.<sup>90</sup> Induction of secretion of interleukin-8 (IL-8) and monocyte chemoattractant protein-1 (MCP-1) has been described in a wide variety of cells while a small group of C-X-C chemokines have only been found preformed and stored in platelet granules. Apart from chemotaxis, chemokines also induce leucocyte activation. IL-8 and MCP-1 induce degranulation and the respiratory burst in neutrophils and monocytes respectively.<sup>103,104</sup> Additionally, while IL-8 induces PAF secretion by neutrophils,<sup>105</sup> MCP-1 induces IL-1 and IL-6 secretion by monocytes.<sup>106</sup>

C-X-C (or  $\alpha$ ) chemokines are so-called because they have an intervening amino acid between the first two of the four conserved cysteines at their N-terminus. They are encoded for on chromosome 4.<sup>107</sup> C-X-C chemokines include IL-8, epithelial-derived neutrophil attractant-78 (ENA-78), growth-related cytokine- $\alpha$  (GRO- $\alpha$ ), interferon  $\gamma$ -inducible protein-10 (IP-10) and platelet factor-4 (PF4). C-X-C chemokines (IL-8, GRO, ENA-78) are primarily neutrophil chemoattractants. However, IP-10 attracts monocytes but not neutrophils<sup>108</sup> while PF4 can attract both neutrophils and monocytes.<sup>109</sup> IL-8 can also attract basophils<sup>110</sup> and by an indirect mechanism, T-cells.

C-C (or  $\beta$ ) chemokines, on the other hand, do not have an intervening amino acid between their first two amino-terminal cysteine residues. They are encoded for on chromosome 17<sup>109</sup> and are primarily monocyte and T-cell chemoattractants.<sup>111,112</sup> C-C chemokines include MCP-1, MCP-2, MCP-3, monocyte inflammatory protein-1 $\alpha$  (MIP-1 $\alpha$ ), MIP-1 $\beta$  and RANTES (regulated on activation, normal T expressed and secreted).

#### *Mechanisms of chemoattractant function*

*In vitro* assays of chemotaxis<sup>113,114</sup> suggest that a "diffusion barrier" located between leucocytes and the "reservoir" of chemoattractant substance is necessary to establish a concentration gradient. Leucocytes sense and crawl "up" this gradient in the direction of increasing concentration. In contrast, leucocytes placed in an environment of uniform chemoattractant concentration undergo enhanced random and directionless movements called "chemokinesis".<sup>115</sup>

*In vivo*, vascular endothelium, the vessel wall, and the interstitial tissue spaces may serve the function of being a "diffusion barrier" that establishes a concentration gradient. Blood flow, by dilution and fluid motion, would actively tend to lower the concentration of chemoattractant in the intravascular compartment. Moreover, red blood cells and venous endothelium bear large numbers of a "scavenger receptor" for chemokines,<sup>116,117</sup> which would tend to lower the concentration of functional chemoattractant within the circulation. However, these factors are absent in the extravascular/interstitial compartment where chemoattractants may accumulate. High levels of IL-8, for example, have been found sequestered in extravascular tissue compartments such as infected joint and pleural cavities.<sup>118–120</sup> IL-8 has been reported to be secreted in a polarised fashion into the extravascular compartment.<sup>121</sup>

In summary, it appears that chemoattractants concentrated in extravascular tissue spaces induce leucocyte recruitment, while high levels of intravascular chemoattractant tend to attenuate leucocyte extravasation.<sup>102,122</sup>




#### **The Leucocyte-endothelial Adhesion Cascade**

The recruitment of leucocytes is precisely regulated by vascular endothelium, which, on activation, presents an adhesive surface to circulating leucocytes. The process of adhesion and diapedesis then involves sequential binding of different CAM to their counter-receptors in a so-called "adhesion cascade". Generally, leucocyte CAM are expressed constitutively, while vascular endothelium express CAM on activation. *In vivo* studies, particularly using intravital microscopy, indicate that leucocyte adhesion and diapedesis mostly occurs in post-capillary venules rather than in arterioles or capillaries.<sup>123,124</sup> This may, in part, be due to higher blood flow rates in arterioles, but may also be related to intrinsic differences in the venular and arteriolar endothelial expression of CAM necessary for adhesion under conditions of flow.

The individual steps of the adhesion cascade became apparent from direct observation of the dynamics of leucocyte-endothelial adhesion, by both intravital videomicroscopy<sup>28,29</sup> as well as by using *in vitro* flow systems in which the pathophysiological and hydrodynamic conditions of leucocyte adhesion under fluid flow at the luminal surface of vascular endothelium were simulated.<sup>27,125</sup> The five main steps of the adhesion cascade are: (1) random contact and margination, (2) initial attachment (or tethering) and rolling, (3) stable arrest and adhesion, (4) transmigration (or diapedesis), (5) interstitial migration and carrying out effector functions (Fig. 1).

#### *Margination (Fig. 1A)*

This first step in leucocyte-endothelial adhesion *in vivo* is a purely random and haemodynamic phenomenon that does not involve the formation of any cell adhesion bonds. Normally, blood flowing through postcapillary venules may be observed to form a central cellular zone consisting of red blood cells and leucocytes and a peripheral cell-free zone.<sup>126</sup> At sites of inflammation, they tend to be pushed away from the centre stream towards the endothelial surface (marginated) by a number of haemodynamic factors.

**A. MARGINATION****Haemodynamic factors****B. ROLLING****Selectins****C. STABLE ADHESION****Ig-integrins / Chemokines****D. DIAPEDESIS/TRANSMIGRATION/CHEMOTAXIS****CD31-CD31 / Ig-integrins / Chemokines****E. INTERSTITIAL MIGRATION/EFFECTOR FUNCTIONS****Chemokines / Ig-integrins**



Smaller-sized red blood cells travelling at greater velocities tend to marginate leucocytes.<sup>127</sup> Leucocytes from smaller tributaries feeding into the side of larger venules are pushed by the prevailing blood flow in the larger vessel towards its sidewalls.<sup>123</sup> During inflammation, acute phase proteins in the circulation increase the viscosity of the blood and may induce the formation of erythrocyte aggregates. This, along with the slowing of blood flow in the microcirculation at sites of inflammation due to vasodilation, will also tend to marginate leucocytes.<sup>128</sup>

The release of vasodilators, e.g. histamine and prostaglandins, during the inflammatory response may increase blood flow and, therefore, the delivery of leucocytes to sites of inflammation. In the presence of tissue injury, upregulated CAM expression may result in adhesive interactions between the marginated leucocytes and the hyperadhesive vascular endothelial surface, thus initiating the adhesion cascade.

#### *Rolling (Figure 1B)*

Leucocytes in postcapillary venules at sites of inflammation may be observed to decelerate and "roll". This is the resultant of transient adhesion bonds being repeatedly formed and broken, between endothelial and leucocyte CAM under the prevailing hydrodynamic forces of fluid flow.

This "rolling" behaviour of leucocytes contacting the luminal surface of activated vascular endothelium is mediated primarily by the selectins,<sup>24,129</sup> although VCAM-1 has also been shown to mediate leucocyte rolling.<sup>130</sup> The shear forces at the luminal surface of postcapillary venules are around 1–5 dynes/cm<sup>2</sup><sup>131,132</sup> and require the formation of selectin-mediated bonds to "capture" marginated leucocytes being swept along by blood flow. At lower flow rates (shear rate  $\leq 0.6$  dynes/cm<sup>2</sup>), L-selectin appears to become less efficient at mediating rolling,<sup>133</sup> and then other adhesion mechanisms, such as  $\beta_2$  integrin-ICAM-1, may function in mediating initial attachment and rolling.<sup>125,134</sup>

Neutrophils have been shown to undergo rolling on both activated vascular endothelium as well as on

other neutrophils already stably adherent to the endothelial surface *in vitro*.<sup>50,135</sup> Neutrophil-neutrophil rolling and aggregation may be a means of amplifying the rate of recruitment of neutrophils at sites of inflammation.<sup>50,136</sup> Free-flowing neutrophils may be seen to tether to activated vascular endothelium and then to form linear aggregates and discrete foci of rolling neutrophils.<sup>27,50</sup> These neutrophil-neutrophil adhesive interactions are mediated by L-selectin binding to PSGL-1. Leucocytes carry both L-selectin as well as PSGL-1 preferentially on the tips of their microvilli, ideally placed for mediating initial attachment and rolling on other neutrophils or vascular endothelium.<sup>46,137</sup>

In summary, a selectin-mediated step is required for leucocyte-endothelial adhesion to occur and culminate in successful leucocyte recruitment, under pathological conditions of fluid flow.

#### *Stable arrest and adhesion (Figure 1C)*

Rolling leucocytes having decelerated begin to form more stable adhesion bonds and, thus, "arrest" in stationary positions on the endothelial surface despite the prevailing shear forces of fluid flow. A necessary prerequisite for this step to occur successfully appears to be integrin activation. Chemokines (e.g. IL-8) and other chemoattractants, such as PAF,<sup>26</sup> are generated at inflammatory sites.<sup>138</sup> Leucocytes rolling on activated endothelium may be exposed to chemokines/chemoattractants or other stimuli which activate the leucocyte integrins rendering them competent to bind ICAM-1 and VCAM-1. Thus, stable arrest in neutrophils and monocytes is mediated by  $\beta_2$  integrin-ICAM-1 adhesion.<sup>27,29</sup> Additionally, mononuclear leucocytes also employ  $\beta_1$  and  $\beta_3$  integrins to bind to VCAM-1<sup>27</sup> and stably adhere to activated vascular endothelium.

#### *Transendothelial migration/diapedesis (Figure 1D)*

Stably adherent leucocytes then begin to flatten and spread on the luminal surface and extend pseudopods

**Fig. 1 (opposite)** (A) Margination is largely a haemodynamic phenomenon and does not depend on any adhesion bonds. (B) Initial attachment and rolling is mediated by selectins (e.g. L-selectin) binding their counter-receptors (e.g. PSGL-1). Neutrophils roll and decelerate on both activated endothelium as well as other adherent neutrophils. (C) Stable adhesion is mediated by adhesion between integrins (e.g. Mac-1 and  $\alpha_4\beta_1$ ) present on the leucocyte surface and Ig (e.g. ICAM-1 and VCAM-1) on the surface of activated endothelium. (D) Transendothelial migration is mediated by CD31-CD31 adhesion between stably adherent leucocytes and endothelium as well as integrin-Ig bonds. Chemokines may promote diapedesis by activating leucocyte integrins or inducing chemotaxis. (E) Chemokines sequestered in extravascular tissue spaces may direct the interstitial migration of transmigrated leucocytes and induce leucocyte activation. Extravasated leucocytes may in turn generate more chemoattractants, thereby amplifying the recruitment process.

towards intercellular borders. This process which is mediated by leucocyte integrins engaging ICAM-1 and/or VCAM-1 appears to involve the coordinated formation and breaking and then re-formation of adhesion bonds as the leucocyte "crawls" across the endothelial surface towards the intercellular border without being swept away by fluid flow. At the intercellular border leucocytes extend pseudopods between endothelial cells and transmigrate into the extravascular spaces. The process of neutrophil and monocyte transmigration or diapedesis between the intercellular junctions of normal and activated vascular endothelium has been shown to be mediated in part by both integrin-ICAM-1 or integrin-VCAM-1 as well as CD31-CD31 adhesive interactions.<sup>25,81,82</sup> The relative importance of each mechanism remains to be further clarified for different leucocyte subsets. For example, some leucocytes (memory T-cells) do not express CD31 and yet are able to transmigrate successfully.<sup>139</sup> Chemoattractants may promote diapedesis by inducing chemotaxis or by integrin activation.

#### *Interstitial migration and effector functions (Figure 1E)*

At sites of inflammation, chemoattractants are generated and then sequester in interstitial and extravascular tissue spaces, being secreted by tissue macrophages and non-vascular cells such as fibroblasts.<sup>119,140-142</sup> Chemokines entering the circulation are carried away by blood flow and become functionally inactive by binding to scavenger receptors found on the surface of red blood cells and venular endothelium.<sup>143</sup> Functionally active chemokines and other chemoattractants thus tend to localise in extravascular tissue compartments and from there induce chemotaxis and activation of extravasated leucocytes, which then are induced to carry out their respective effector functions such as cytokine secretion, phagocytosis and degranulation.

### **Mechanisms of Amplifying the Inflammatory Response**

A number of mechanisms exist which seem to both regulate and amplify the inflammatory response. L-selectin-PSGL-1-mediated adhesion and aggregation between neutrophils may be a means of localising and amplifying the numbers of neutrophils "captured" to adhere to activated endothelium at sites of inflammation.<sup>50</sup> Inflammatory mediators may be sequentially generated by different cells. For example,

C5a produced during myocardial ischaemia-reperfusion injury mediates recruitment of neutrophil which in turn, on activation, may generate IL-8, PAF and LTB<sub>4</sub>, thus inducing further neutrophil recruitment.<sup>144,145</sup> Similar mechanisms may exist for monocytes and other leucocytes.<sup>146</sup> Furthermore, there appears to be a system of networking between different cells via cytokines and other chemoattractants to amplify the inflammatory response. During Gram-negative pulmonary sepsis, alveolar macrophages, on activation by endotoxin, secrete inflammatory cytokines which then act on pulmonary fibroblasts to generate IL-8, while endotoxin alone did not induce significant IL-8 secretion by pulmonary fibroblasts.<sup>141</sup>

### **Endothelial-mediated Mechanisms of Atherosclerosis**

The importance of atherosclerosis as a leading cause of morbidity and mortality is undisputed. Atherosclerosis is thought to be an excessive inflammatory-fibroproliferative response to chronic injury in susceptible arteries.<sup>147</sup> It appears to be the result of complex interactions between arterial endothelium and smooth muscle cells, and surrounding physical and biochemical factors such as circulating leucocytes and platelets, proteins and lipids carried in plasma and haemodynamic forces.<sup>2</sup>

The earliest anatomical lesion of atherosclerosis is a subendothelial accumulation of lipid-laden macrophages called the "fatty streak". An even earlier cellular event during atherogenesis is the increased adherence of peripheral blood monocytes to the endothelium of large arteries at sites which later develop fatty streaks.<sup>148</sup> In a rabbit model of atherogenesis, recruitment of peripheral blood monocytes is preceded by focal VCAM-1 expression.<sup>149</sup> VCAM-1, expressed by activated arterial endothelium, mediates monocyte-endothelial adhesion,<sup>27</sup> and may be one of the earliest molecular markers of atherosclerosis. At these sites, fatty streaks develop and later progress into fibro-fatty atherosclerosis plaques containing monocytes and T-lymphocytes.<sup>150</sup> Other CAM, namely, ICAM-1, CD31, and P-selectin, have also been shown to be expressed by human atherosclerotic plaques<sup>151,152</sup> and each of these is capable of mediating leucocyte-endothelial adhesion.<sup>27,82,153</sup>

Hyperlipidaemia *per se* results in a number of pro-atherogenic changes in the vessel wall. It induces endothelial activation and the expression of MCP-1, a monocyte chemoattractant.<sup>154</sup> Hyperlipidaemia also stimulates monocyte production of IL-1 $\beta$ ,<sup>155,156</sup> which in

turn induces further endothelial activation and MCP-1 expression. IL-1 $\beta$  may also act as a growth factor for smooth muscle cells.<sup>157</sup> Furthermore, cytokine activation of endothelium and smooth muscle cells has been shown to induce the expression of macrophage colony stimulating factor,<sup>158</sup> a cytokine involved in the growth and maturation of monocyte-macrophages. Diabetics develop premature and florid atherosclerosis and a possible mechanism for this may involve the formation of advanced glycation endproducts in the vascular wall in response to chronic hyperglycaemia. These advanced glycation endproducts have been shown to promote monocyte adhesion and chemotaxis.<sup>159,160</sup> Monocyte-macrophages, which are present at all stages of atherogenesis, may in turn release a number of inflammatory cytokines and growth factors involved in the progression of the atherosclerotic plaque.<sup>159</sup>

### Therapeutic Applications

While endothelial-leucocyte adhesion and diapedesis are necessary for normal host defence, there exist a number of inflammatory diseases in which excessive or inappropriate leucocyte recruitment contributes to tissue damage. For example, in atherosclerosis, mononuclear leucocytes accumulate in the arterial wall and secrete inflammatory cytokines and growth factors which promote plaque progression. Although the pathogenesis of atherosclerosis is not fully understood, it is believed to be an inflammatory response to injurious stimuli.<sup>159</sup> Therefore, the ability to inhibit the recruitment of mononuclear leucocytes may be a way of modulating plaque progression. Similarly, restoration of blood flow following a period of limb or organ ischaemia is characterised by recruitment and activation of neutrophils which are believed to induce postischaemic tissue injury during reperfusion. Platelet thrombosis occurs by two cell adhesion processes (platelet adherence and aggregation), both of which are mediated by platelet integrins and glycoproteins. All these disease processes depend on the adhesion and aggregation of leucocytes and platelets to vascular endothelium mediated by CAM and chemoattractants.

### Potential molecular targets

Knowledge of the cellular and molecular mechanisms of leucocyte recruitment and platelet thrombosis affords the possibility of pharmacologically modulating

these cell adhesion processes and a number of molecular sites present themselves as being potential targets (Table 3). Antagonism of the cytokines which drive the inflammatory response may reduce the severity of tissue damage locally and dampen undesirable systemic effects. Disruption of some or all of the steps of the endothelial-leucocyte adhesion cascade, by blockade of CAM and/or chemoattractants, may be a means of controlling leucocyte recruitment in inflammatory diseases. Instead of blockade at the level of extracellular protein, it may be possible to antagonise the expression of one or more inflammatory response genes at the cytoplasmic transcription factor or proteasomal level.<sup>161,162</sup> Finally, endothelial-leucocyte adhesion may be modulated by reducing the systemic leucocyte count by the use of extracorporeal leucocyte filters.

### Possible limitations

The temporal and spatial patterns of expression of CAM *in vivo* in various disease states have yet to be fully characterised, and, therefore, it may not be possible at present to effectively block a particular CAM in a specific and timely fashion. Furthermore, the existence of multiple or redundant mechanisms of leucocyte recruitment in particular organs and diseases may necessitate the simultaneous blockade of multiple adhesion pathways. For example, in the pulmonary microcirculation, CD18-dependent and CD18-independent mechanisms of leucocyte recruitment appear to operate.<sup>163</sup>

Monoclonal antibody (mAb) therapy has certain theoretical limitations. mAb generated using animals may induce sensitisation or contain endotoxin or virus contamination. The Fc portion of mAb may mediate complement activation and non-specific neutrophil binding once the F(ab')<sub>2</sub> portion has bound to the endothelial target antigen. However, while F(ab')<sub>2</sub> fragments are able to bind and block the target antigen, they have brief plasma half-lives being cleared relatively rapidly from the circulation. Additionally, in animal studies, mAb have been shown not to be efficacious in reducing leucocyte recruitment unless their efficiency of blockade exceeded 90%.<sup>164</sup> Once a few neutrophils adhere, these appear able to recruit other neutrophils, presumably by amplification mechanisms (see above).

A theoretical concern of blockade of  $\beta_2$  (CD18) integrins is the possibility of inducing an immunocompromised state and uncontrolled sepsis by interfering with the ability of neutrophils to extravasate and kill bacteria at sites of infection. Animal

**Table 3. Molecular targets in the modulation of inflammation and thrombosis.**

Target molecule	Function	Pharmacological agent
Inflammatory cytokines	Endothelial and leucocyte activation	Monoclonal antibody Soluble receptor antagonist
Selectins	Leucocyte-endothelial initial attachment and rolling	Monoclonal antibody Soluble selectin or ligand Synthetic carbohydrate
$\beta_2$ integrins Ig	Stable adhesion of leucocytes and diapedesis	Monoclonal antibody Soluble CAM Soluble counter-receptor
gpIIb/IIIa Platelet integrins	Platelet aggregation Platelet-endothelial adhesion	Monoclonal antibody
Chemoattractants Chemokines	Integrin activation and leucocyte recruitment	Monoclonal antibody Soluble receptor Synthetic receptor antagonist
Transcription regulation of inflammatory response genes	mRNA synthesis and protein synthesis of inflammatory response gene product e.g. E-selectin	Proteasome inhibitors Antisense oligonucleotides

studies have yielded conflicting results as to whether CD18-blockade increased septic complications.<sup>165,166</sup>

#### *Ischaemia-reperfusion injury*

Tissue ischaemia may occur in a variety of clinical situations including acute vessel occlusion by thromboembolism, trauma or surgical cross-clamping of major vessels, compartment syndrome, autogenous muscle transplantation for wound coverage and solid organ transplantation. During ischaemia, cellular injury is caused by a reduction in the supply of oxygen and nutrients as well as inadequate clearance of products of metabolism. Survival of ischaemic tissue is, therefore, dependent on timely and sufficient restoration of its blood supply (reperfusion). However, beyond a certain period of ischaemia, reperfusion may itself result in additional damage of postischaemic tissues and an accelerated rate of necrosis despite adequate restoration of blood flow.<sup>167,168</sup> Neutrophil infiltration is characteristic of the inflammatory response to reperfusion following ischaemia.<sup>168,169</sup> The pathogenesis of reperfusion injury is complex but is believed to be due in part to neutrophil-mediated damage.<sup>168,170–172</sup> Both CAM and chemoattractants have been shown to mediate neutrophil recruitment in reperfusion injury.<sup>144,164,172</sup> Blockade of CAM and chemoattractants in numerous animal models of ischaemia and reperfusion in different organs appeared to reduce the extent of tissue damage during reperfusion.<sup>144,164,168,172</sup> These findings remain to be confirmed in clinical studies.

#### *Platelet thrombosis, restenosis and thrombolysis*

The surface exposed by vascular endothelial injury following balloon angioplasty or ulceration of atherosclerotic plaque is highly thrombogenic. Platelets adhere to subendothelial matrix proteins such as collagen and fibronectin, and then become activated and degranulate, releasing coagulation proteins, calcium ions (necessary for coagulation), ADP (mediates platelet aggregation), and chemokines (e.g. PF4). Activated platelets may also secrete thromboxane  $A_2$ , a potent vasoconstrictor and stimulus of platelet aggregation.

Low-dose aspirin inhibits the synthesis of thromboxane  $A_2$  by irreversibly acetylating the enzyme cyclooxygenase. As platelets are anuclear cells unable to synthesise new proteins, they cannot manufacture new enzyme during their short lifespan of just over a week. In contrast, prostacyclin (an inhibitor of platelet aggregation) production by endothelial cells is relatively unaffected by low doses of aspirin. Thus, low dose aspirin tips the balance of prostaglandin metabolism towards inhibition of platelet aggregation.<sup>173</sup> Ticlopidine and clopidogrel are novel antiplatelet agents which are more potent than aspirin.<sup>174</sup> They are inactive *in vitro*, but *in vivo*, after hepatic metabolism into their active moieties, they produce irreversible platelet inhibition by selectively blocking ADP receptors and inhibiting fibrinogen binding of gpIIb/IIIa.<sup>175</sup>

Following balloon angioplasty of atherosclerotic vessels, damage to the intima may induce platelet activation and aggregation and thrombosis resulting in acute thrombotic occlusion (termed "abrupt closure"). Activated platelets may also be a source of growth



factors which mediate smooth muscle cell proliferation that characterises both progression of atherosclerotic plaques as well as late restenosis of angioplastied lesions. A chimeric mAb F(ab')<sub>2</sub> fragment against gpIIb/IIIa (c7E3; trade name ReoPro) has been shown in clinical trials to reduce the incidence of acute ischaemic complications due to platelet thrombosis following coronary angioplasty and atherectomy.<sup>176,177</sup> Unfortunately, the rate of restenosis has not been consistently reduced. However, blockade of platelet thrombosis with anti-gpIIb/IIIa mAb following coronary angioplasty significantly reduced the rates of death, myocardial infarction and need for further revascularisation.<sup>178</sup> Blockade of platelet thrombosis may also prove to be useful in selected cases of peripheral e.g. infrainguinal angioplasty.

It should be noted that platelets appear to have both profibrinolytic as well as antifibrinolytic activity. Both streptokinase and tissue-plasminogen activator induce platelet activation,<sup>179</sup> and platelet-rich thrombi appear to be more resistant to lysis.<sup>180</sup> On the other hand, thrombolysis may be accelerated in the presence of platelets by binding plasminogen via a gpIIb/IIIa-dependent mechanism, thus generating high local concentrations of plasmin.<sup>181</sup> The concomitant use of antiplatelet agents during thrombolysis seems to expedite reperfusion and may lower the dose of thrombolytic agent required to attain reperfusion, and so reduce the risk of haemorrhage.<sup>86</sup> Further studies are necessary to determine the effects of platelets and antiplatelet therapy during thrombolysis in different clinical settings.

## Conclusions

Thus, the pathogenesis of a number of vascular diseases and their complications depend on cell adhesion processes mediated by CAM and chemoattractants and occur as vascular endothelial cells respond to injurious stimuli. The continued understanding of these molecular processes will allow the development of therapeutic strategies in the future to treat and arrest the progression of these diseases by modulating leucocyte-endothelial and platelet-endothelial adhesion.

## References

- HUNTER J. Treatise on the blood inflammation and gunshot wounds. In: Palmer JF, ed. *The Surgical Works of John Hunter*, F.R.S. London: Longman, Rees, Orme, Brown, Green and Longman, 1835: 1-580.
- GIMBROKE JR MA, CYBULSKY MI, KUME N, COLLINS T, RESNICK N. Vascular endothelium. An integrator of pathophysiological stimuli in atherogenesis. *Ann New York Acad Sci* 1995; **748**: 122-132.
- POBER JS, COTRAN RS. Overview: the role of endothelial cells in inflammation. *Transplantation* 1990; **50**: 537-544.
- FRANGOS JA, ESKIN SG, MCINTYRE LV, IVES CL. Flow effects on prostacyclin production by cultured human endothelial cells. *Science* 1985; **227**: 1477-1479.
- RUBANYI GM, ROMERO JC, VANHOUTTE PM. Flow induced release of endothelium-derived relaxing factor. *Am J Physiol* 1986; **250**: H1145-H1149.
- NAGEL T, RESNICK N, ATKINSON WJ, DEWEY JR CF, GIMBRONE JR MA. Shear stress selectively up-regulates intercellular adhesion molecule-1 expression in cultured human vascular endothelial cells. *J Clin Invest* 1994; **94**: 885-891.
- LUSCINSKAS FW, GIMBRONE JR MA. Endothelial-dependent mechanisms in chronic inflammatory leukocyte recruitment. *Annu Rev Med* 1996; **47**: 413-421.
- WELLICOME SM, THORNHILL MH, PITZALIS C *et al*. A monoclonal antibody that detects a novel antigen on endothelial cells that is induced by tumor necrosis factor, IL-1 or lipopolysaccharide. *J Immunol* 1990; **144**: 2558-2565.
- MCIVER RP, BECKSTEAD JH, MOORE KL, MARSHALL-COLSON L, BANTON DF. GMP-140, a platelet alpha-granule membrane protein, is also synthesized by vascular endothelial cells and is localized in Weibel-Palade bodies. *J Clin Invest* 1989; **84**: 92-106.
- BEVILACQUA MP, POBER JS, MENDRICK DL, COTRAN RS, GIMBRONE JR MA. Identification of an inducible endothelial-leukocyte adhesion molecule. *Proc Natl Acad Sci USA* 1987; **84**: 9238-9242.
- POBER JS, GIMBRONE JR MA, LAPIERRE LA *et al*. Overlapping patterns of activation by human endothelial cells by interleukin-1, tumor necrosis factor and immune interferon. *J Immunol* 1986; **137**: 1893-1896.
- MCINTYRE TM, ZIMMERMAN GA, PRESCOTT SM. Leukotrienes C<sub>4</sub> and D<sub>4</sub> stimulate human endothelial cells to synthesize platelet-activating factor and bind neutrophils. *Proc Natl Acad Sci USA* 1986; **83**: 2204-2208.
- GIMBRONE JR MA, OBIN MS, BROCK AF *et al*. Endothelial interleukin-8: A novel inhibitor of leukocyte-endothelial interactions. *Science* 1989; **246**: 1601-1603.
- COLLINS T. Biology of disease. Endothelial nuclear factor-κB and the initiation of the atherosclerotic lesion. *Lab Invest* 1993; **68**: 499-508.
- BEVILACQUA MP, STENGELIN S, GIMBRONE JR MA, SEED B. Endothelial leukocyte adhesion molecule 1: an inducible receptor for neutrophils related to complement regulatory proteins and lectins. *Science* 1989; **243**: 1160-1165.
- ESMON CT. The protein C anticoagulant pathway. *Arterioscler Thromb* 1992; **12**: 135-145.
- CARMELIET P, COLLEN D. Gene targeting and gene transfer studies of the plasminogen/plasmin system: implications in thrombosis, hemostasis, neo-intima formation and atherosclerosis. *FASEB J* 1995; **9**: 934-938.
- MACKMAN N. Regulation of the tissue factor gene. *FASEB J* 1995; **9**: 883-889.
- MARTIN DMA, BOYS CWG, RUF W. Tissue factor: molecular recognition and co-factor function. *FASEB J* 1995; **9**: 852-859.
- SAWA H, FUJII S, SOBEL BE. Augmented arterial wall expression of type-1 plasminogen activator inhibitor induced by thrombosis. *Arterioscler Thromb* 1992; **12**: 1507-1515.
- METCHNIKOFF E. *Lectures on the Comparative Pathology of Inflammation*. New York: Dover Publications Inc., (re-publication of original edition by Kegan, Paul, Trench Trubner and Co., London, 1893), 1968.
- LARSON RS, SPRINGER TA. Structure and function of leukocyte integrins. *Immunol Rev* 1990; **114**: 181-217.
- CARLOS TM, HARLAN JM. Leukocyte-endothelial adhesion molecules. *Blood* 1994; **84**: 2068-2101.



- 24 KANSAS GS. Selectins and their ligands: current concepts and controversies. *Blood* 1996; **88**: 3259–3287.
- 25 LUSCINSKAS FW, CYBULSKY MI, KIELY JM *et al*. Cytokine-activated human endothelial monolayers support enhanced neutrophil transmigration via a mechanism involving both endothelial-leukocyte adhesion molecule-1 and intercellular adhesion molecule-1. *J Immunol* 1991; **146**: 1617–1625.
- 26 LORANT DE, PATEL KD, MCINTYRE TM *et al*. Coexpression of GMP-140 and PAF by endothelium stimulated by histamine or thrombin: a juxtacrine system for adhesion and activation of neutrophils. *J Cell Biol* 1991; **115**: 223–234.
- 27 LUSCINSKAS FW, KANSAS GS, DING H *et al*. Monocyte rolling, arrest and spreading on IL-4-activated vascular endothelium under flow is mediated via sequential action of L-selectin,  $\beta_1$ -integrins and  $\beta_2$ -integrins. *J Cell Biol* 1994; **125**: 1417–1427.
- 28 LEY K, GAEHTGENS P, FENNIE C *et al*. Lectin-like cell adhesion molecule-1 mediates rolling in mesenteric venules in vivo. *Blood* 1991; **77**: 2553–2555.
- 29 VON ANDRIAN UH, CHAMBERS JD, McEVOY LM *et al*. Two-step model of leukocyte-endothelial cell interactions in inflammation: distinct roles for LECAM-1 and the leukocyte  $\beta_2$  integrins in vivo. *Proc Natl Acad Sci USA* 1991; **88**: 7538–7542.
- 30 MAYADAS TN, JOHNSON RC, RAYBURN H, HYNES RO, WAGNER DD. Leukocyte rolling and extravasation are severely compromised in P-selectin-deficient mice. *Cell* 1993; **74**: 541–554.
- 31 ARBONES MI, ORD DC, LEY K *et al*. Lymphocyte homing and leukocyte rolling and migration are impaired in L-selectin (CD62L)-deficient mice. *Immunity* 1994; **1**: 247–260.
- 32 LABOW MA, NORTON CR, RUMBERGER JM *et al*. Characterization of E-selectin deficient mice demonstration of overlapping function of the endothelial selectins. *Immunity* 1994; **1**: 709–720.
- 33 BULLARD DC, QIN L, LORENZO L *et al*. P-selectin/ICAM-1 double mutant mice acute emigration of neutrophils into the peritoneum is completely absent but is normal into pulmonary alveoli. *J Clin Invest* 1995; **95**: 1782–1788.
- 34 ARNAOUT MA. Leukocyte adhesion molecule deficiency, its structural basis, pathophysiology and implications for modulating the inflammatory response. *Immunol Rev* 1990; **114**: 147–180.
- 35 PRICE TH, OCHS HD, GERSHONI-BARUCH R, HARLAN JM, ETZIONI A. In vivo neutrophil and lymphocyte function studies in a patient with leukocyte adhesion deficiency type II. *Blood* 1994; **84**: 1635–1639.
- 36 SPRINGER TA. Traffic signals for lymphocyte recirculation and leukocyte emigration: the multistep paradigm. *Cell* 1994; **76**: 301–314.
- 37 TEDDER TF, LUSCINKAS FW, KANSAS GS. Regulation of leukocyte migration by L-selectin: mechanism, domains and ligands. *Behring Inst Mitt* 1993; **92**: 165–177.
- 38 KISHIMOTO TK, JUTILA MA, BERG EL, BUTCHER EC. Neutrophil Mac-1 and MEL-14 adhesion proteins inversely regulated by chemotactic factors. *Science* 1989; **245**: 1238–1241.
- 39 GEARING AJ, NEWMAN W. Circulating adhesion molecules in disease. *Immunol Today* 1994; **14**: 506–512.
- 40 SCHLEIFFENBAUM B, SPERTINI O, TEDDER TF, Soluble L-selectin is present in human plasma at high levels and retains functional activity. *J Cell Biol* 1992; **119**: 229–238.
- 41 ALLPORT JR, DING H, AGER A *et al*. L-selectin shedding does not regulate human neutrophil attachment, rolling, or transmigration across human vascular endothelium. *J Immunol* 1997; **158**: 4368–4372.
- 42 KOCH AE, BURROWS JC, HAINES GK *et al*. Immunolocalization of endothelial and leukocyte adhesion molecules in human rheumatoid and osteoarthritic synovial tissues. *Lab Invest* 1991; **64**: 313–320.
- 43 PICKER LJ, KISHIMOTO TK, SMITH CW, WARNOCK RA, BUTCHER EC. ELAM-1 is an adhesion molecule for skin-homing T-cells. *Nature* 1991; **349**: 796–799.
- 44 VARKI A. Selectin ligands. *Proc Natl Acad USA* 1994; **91**: 7390–7397.
- 45 SAKO D, CHANG XJ, BARONE KM *et al*. Expression cloning of a functional glycoprotein ligand for P-selectin. *Cell* 1993; **75**: 1179–1186.
- 46 MOORE KL, PATEL KD, BRUEHL RE *et al*. P-selectin glycoprotein ligand-1 mediates rolling of human neutrophils on P-selectin. *J Cell Biol* 1995; **126**: 661–671.
- 47 GUYER DA, MOORE KL, LYNAM EB *et al*. P-selectin glycoprotein ligand (PSGL-1) is a ligand for L-selectin in neutrophil aggregation. *Blood* 1996; **88**: 2415–2421.
- 48 GOETZ DJ, GREIF DM, DING H *et al*. Isolated P-selectin glycoprotein ligand-1 dynamic adhesion to P- and E-selectin. *J Cell Biol* 1997; **137**: 509–519.
- 49 PATEL KD, MOORE KL, NOLLERT MU, McEVER RP. Neutrophils use both shared and distinct mechanisms to adhere to selectins under static and flow conditions. *J Clin Invest* 1996; **96**: 1887–1896.
- 50 WALCHECK B, MOORE KL, McEVER RP, KISHIMOTO TK. Neutrophil-neutrophil interactions under hydrodynamic shear stress involve L-selectin and PSGL-1. A mechanism that amplifies initial leukocyte accumulation of P-selectin in vitro. *J Clin Invest* 1996; **98**: 1081–1087.
- 51 SPERTINI O, LUSCINSKAS FW, KANSAS GS *et al*. Leukocyte adhesion molecule-1 (LAM-1, L-selectin) interacts with an inducible endothelial cell ligand to support leukocyte adhesion. *J Immunol* 1991; **147**: 2565–2573.
- 52 LUSCINSKAS FW, LAWLER J. Integrins as dynamic regulators of vascular function. *FASEB J* 1994; **8**: 929–938.
- 53 ANDERSON DC, MILLER LJ, SCHMALSTIEG FC, ROTHLEIN R, SPRINGER TA. Contributions of the Mac-1 glycoprotein family to adherence-dependent granulocyte functions structure-function assessment employing subunit-specific monoclonal antibodies. *J Immunol* 1986; **137**: 15–27.
- 54 BAINTON DF, MILLER LJ, KISHIMOTO TK, SPRINGER TA. Leukocyte adhesion receptors are stored in peroxidase-negative granules of human neutrophils. *J Exp Med* 1987; **166**: 1641–1653.
- 55 DIAMOND MS, SPRINGER TA. A subpopulation of Mac-1 (CD11b/CD18) molecules mediates neutrophil adhesion to ICAM-1 and fibrinogen. *J Cell Biol* 1993; **120**: 545–566.
- 56 FEARON DT, COLLINS LA. Increased expression of C3b receptors on polymorphonuclear leukocytes induced by chemotactic factors and by purification procedures. *J Immunol* 1983; **130**: 370–375.
- 57 ARNAOUT MA. Structure and function of leukocyte adhesion molecules CD11/CD18. *Blood* 1990; **75**: 1037–1050.
- 58 DRANSFIELD I, HOGG N. Regulated expression of Mg<sup>2+</sup> binding epitope on leukocyte integrin  $\alpha$  subunits. *EMBO J* 1989; **8**: 3759–3765.
- 59 DUSTIN ML, SPRINGER TA. Lymphocyte function associated antigen-1 (LFA-1) interaction with intercellular adhesion molecule-1 (ICAM-1) is one of at least three mechanisms for lymphocyte adhesion to cultured endothelial cells. *J Cell Biol* 1988; **107**: 321–331.
- 60 STAUNTON DE, DUSTIN ML, SPRINGER TA. Functional cloning of ICAM-2, a cell adhesion ligand for LFA-1 homologous to ICAM-1. *Nature* 1989; **339**: 61–64.
- 61 DE FOUGEROLLES AR, SPRINGER TA. Intercellular adhesion molecule 3, a third adhesion counter-receptor for lymphocyte function associated molecule-1 on resting lymphocytes. *J Exp Med* 1992; **175**: 185–190.
- 62 DIAMOND MS, STAUNTON DE, MARLIN SD, SPRINGER TA. Binding of the integrin Mac-1 (CD11b/CD18) to the third Ig-like domain of ICAM-1 (CD54) and its regulation by glycosylation. *Cell* 1991; **65**: 961–971.
- 63 BELLER DI, SPRINGER TA, SCHREIBER RD. Anti-Mac-1 selectively inhibits the mouse and human type 3 complement receptor. *J Exp Med* 1982; **156**: 1000–1009.
- 64 MYONES BL, DALZELL JG, HOGG N, ROSS GD. Neutrophil and monocyte cell surface p150,95 has iC3b-receptor (CR4) activity resembling CR3. *J Clin Invest* 1988; **82**: 640–651.
- 65 ROTHLEIN R, DUSTIN ML, MARLIN SD, SPRINGER TA. A human intercellular adhesion molecule (ICAM-1) distinct from LFA-1. *J Immunol* 1986; **137**: 1270–1274.

- 66 MARLIN SD, STAUNTON DE, SPRINGER TA *et al.* A soluble form of intercellular adhesion molecule-1 (ICAM-1) inhibits rhinovirus infection. *Nature* 1990; **344**: 70–72.
- 67 BERENDT AR, SIMMONS DL, TANSY J, NEWBOLD CI. Intercellular adhesion molecule-1 is an endothelial cell adhesion receptor for *Plasmodium falciparum*. *Nature* 1989; **341**: 57–59.
- 68 MEYER DM, DUSTIN ML, CARRON CP. Characterization of intercellular adhesion molecule-1 ectodomain (sICAM-1) as an inhibitor of lymphocyte function-associated molecule-1 interaction with ICAM-1. *J Immunol* 1989; **155**: 3578–3584.
- 69 DE FOUGEROLLES AR, STACKER SA, SCHWARTING R, SPRINGER TA. Characterization of ICAM-2 and evidence for a third counter-receptor for LFA-1. *J Exp Med* 1991; **174**: 253–267.
- 70 RICE GE, MUNRO JM, BEVILACQUA MP. Inducible cell adhesion molecule 110 (INCAM-110) is an endothelial receptor for lymphocytes. A CD11/CD18 independent mechanism. *J Exp Med* 1990; **171**: 1369–1374.
- 71 OSBORN L, HESSION C, TIZARD R *et al.* Direct expression cloning of vascular adhesion molecule-1, a cytokine induced endothelial protein that binds to lymphocytes. *Cell* 1989; **59**: 1203–1211.
- 72 MARLOR CW, WEBB DL, BOMBARA MP, GREVE JM, BLUE ML. Expression of vascular cell adhesion molecule-1 in fibroblast-like synoviocytes after stimulation with tumor necrosis factor. *Am J Pathol* 1992; **140**: 1055–1060.
- 73 BOCHNER BS, LUSCINSKAS FW, GIMBRONE JR MA *et al.* Adhesion of human basophils, eosinophils and neutrophils to interleukin-1 activated human vascular endothelial cells: contributions of endothelial cell adhesion molecules. *J Exp Med* 1991; **173**: 1553–1556.
- 74 KUME N, CYBULSKY MI, GIMBRONE JR MA. Lysophosphatidylcholine, a component of atherogenic lipoproteins, induces mononuclear leukocyte adhesion molecules in cultured arterial endothelial cells. *J Clin Invest* 1992; **90**: 1138–1144.
- 75 SCHLEIMER RP, STEBINSKY SA, KAISER J *et al.* Interleukin-4 induces adherence of human eosinophils and basophils but not neutrophils to endothelium. *J Immunol* 1992; **148**: 1086–1092.
- 76 ELICES MJ, OSBORN L, TANAKA Y *et al.* VCAM-1 on activated endothelium interacts with the leukocyte integrin VLA-4 at a binding site distinct from the VLA-4/fibronectin binding site. *Cell* 1990; **60**: 577–584.
- 77 BERLIN C, BARGATZE RF, CAMPBELL JJ *et al.* Alpha 4 integrins mediate lymphocyte attachment and rolling under physiologic flow. *Cell* 1995; **80**: 413–422.
- 78 ALBELDA SM, MULLER WA, BUCK CA, NEWMAN PJ. Molecular and cellular properties of PECAM-1 (EndoCAM/CD31). A novel vascular cell-cell adhesion molecule. *J Cell Biol* 1991; **114**: 1059–1068.
- 79 ROMER LH, MCLEAN NV, YAN HC *et al.* IFN-gamma and TNF-alpha induce redistribution of PECAM-1 (CD31) on human endothelial cells. *J Immunol* 1995; **154**: 6582–6592.
- 80 RIVAL Y, DEL MASCHIO A, RABET MJ, DEJANA E, DUPERRAY A. Inhibition of platelet endothelial cell adhesion molecule-1 synthesis and leukocyte transmigration in endothelial cells by the combined action of TNF-alpha and IFN-gamma. *J Immunol* 1996; **157**: 1233–1241.
- 81 MULLER WA, WEIGL SA, DENG X, PHILLIPS DM. PECAM-1 is required for transendothelial migration of leukocytes. *J Exp Med* 1993; **178**: 449–460.
- 82 LIAO F, HUYNH HK, EIROA A *et al.* Migration of monocytes across endothelium and passage through extracellular matrix involve separate molecular domains of PECAM-1. *J Exp Med* 1996; **182**: 1337–1343.
- 83 PIALI L, HAMMEL P, UHEREK C *et al.* CD31/PECAM-1 is a ligand for  $\alpha_5\beta_1$  integrin involved in adhesion of leukocytes to endothelium. *J Cell Biol* 1995; **130**: 451–460.
- 84 PALABRICA T, LOBB R, FURIE BC *et al.* Leukocyte accumulation promoting fibrin deposition is mediated in vivo by P-selectin on adherent platelets. *Nature* 1992; **359**: 848–851.
- 85 KIEFFER N, PHILLIPS DR. Platelet membrane glycoproteins: Functions in cellular interactions. *Annu Rev Cell Biol* 1990; **6**: 329–357.
- 86 COLLIER BS. Platelets and thrombolytic therapy. *N Engl J Med* 1990; **322**: 33–42.
- 87 PHILLIPS DR, CHARO IF, PARISE LV, FITZGERALD LA. The platelet membrane glycoprotein IIb-IIIa complex. *Blood* 1988; **71**: 831–843.
- 88 FLOW EF, GINSBERG MH. Cellular adhesion gpIIb-IIIa as a prototype adhesion receptor. *Prog Hemost Thromb* 1989; **9**: 117–156.
- 89 MOVAT HZ. Chemotaxis. In: Movat HZ, ed. *The Inflammatory Reaction*. Amsterdam: Elsevier, 1985: 203–257.
- 90 OPPENHEIM JJ, ZACHARIAE COC, MUKAIDA N, MATSUSHIMA K. Properties of the novel proinflammatory supergene “intercrine” cytokine family. *Ann Rev Immunol* 1991; **9**: 617–648.
- 91 BEN-BARUCH A, MICHIEL DF, OPPENHEIM JJ. Signals and receptors involved in recruitment of inflammatory cells. *J Biol Chem* 1995; **270**: 11703–11706.
- 92 GERARD C, GERARD NP. Molecular biology of human neutrophil chemotactic receptors. In: Hellewell PG, Williams TJ, eds. *Immunopharmacology of Neutrophils*. London: Academic Press, 1994: 115–131.
- 93 KELVIN DJ, MICHIEL DF, JOHNSTON JA *et al.* Chemokines and serpentine: the molecular biology of chemokine receptors. *J Leuk Biol* 1993; **54**: 604–612.
- 94 DEVREOTES PN, ZIGMOND SH. Chemotaxis in eukaryotic cells: a focus on leukocytes and Dictyostelium. *Ann Rev Cell Biol* 1988; **4**: 649–686.
- 95 GOETZL EJ, PICKET WC. Novel structural determinants of the human neutrophil chemotactic activity of leukotriene B. *J Exp Med* 1981; **153**: 482–487.
- 96 MCMAUS LM, WOODARD DS, DEEVERS SI, PINCKARD RN. PAF molecular heterogeneity: pathobiological implications. *Lab Invest* 1993; **69**: 639–650.
- 97 PINCKARD RN, JACKSON EM, HOPPENS C *et al.* Molecular heterogeneity of platelet-activating factor produced by stimulated human polymorphonuclear leukocytes. *Biochem Biophys Res Commun* 1984; **122**: 325–332.
- 98 PRESCOTT SM, ZIMMERMAN GA, MCINTYRE TM. Human endothelial cells in culture produce platelet-activating factor (1-alkyl-2-acetyl-sn-glycero-3-phosphocholine) when stimulated with thrombin. *Proc Nat Acad Sci USA* 1984; **81**: 3534–3538.
- 99 EVANGELOU AM. Platelet-activating factor (PAF): implications for coronary heart and vascular diseases. *Prostaglandins Leukotrienes and Essential Fatty Acids* 1994; **50**: 1–28.
- 100 WARD PA, NEWMAN LJ. A neutrophil chemotactic factor from human C5. *J Immunol* 1969; **102**: 93–99.
- 101 SCHIFFMAN E, CORCORAN B, WAHL SM. N-formylmethionyl peptides as chemoattractants for leukocytes. *Proc Nat Acad Sci USA* 1975; **72**: 1059–1062.
- 102 FURIE MB, RANDOLPH GJ. Chemokines and tissue injury. *Am J Pathol* 1995; **146**: 1287–1301.
- 103 DJEU JY, MATSUSHIMA K, OPPENHEIM JJ, SHIOTSUKI K, BLANCHARD DK. Functional activation of human neutrophils by recombinant monocyte-derived neutrophil chemotactic factor. *J Immunol* 1990; **144**: 2205–2210.
- 104 ZACHARIAE COC, ANDERSON AO, THOMPSON HL *et al.* Properties of monocyte chemotactic and activating factor purified from a human fibrosarcoma cell line. *J Exp Med* 1990; **171**: 2177–2182.
- 105 BUSSOLINO F, SIRONI M, BOCCHIETTO E, MANTOVANI A. Synthesis of platelet-activating factor by polymorphonuclear neutrophils stimulated with IL-8. *J Biol Chem* 1992; **267**: 14598–14603.
- 106 JIANG Y, BELLER DI, FRENDL G, GRAVES DT. Monocyte chemoattractant protein-1 regulates adhesion molecule expression and cytokine production in human monocytes. *J Immunol* 1992; **148**: 2423–2428.
- 107 MODI WS, DEAN M, SEUANEZ HN *et al.* Monocyte-derived neutrophil chemotactic factor (MDNCF/IL-8) resides in a gene cluster along with several other members of the platelet factor gene family. *Hum Genet* 1990; **84**: 185–187.
- 108 TAUB DD, LLOYD AR, CONLON K *et al.* Recombinant human

- interferon-inducible protein 10 is a chemoattractant for human monocytes and T lymphocytes and promotes T cell adhesion to endothelial cells. *J Exp Med* 1993; **177**: 1809–1814.
- 109 DEUEL TF, SENIOR RM, CHAND D *et al*. Platelet factor 4 is chemotactic for neutrophils and monocytes. *Proc Natl Acad Sci USA* 1981; **78**: 4584–4587.
  - 110 LEONARD EJ, SKEEL A, YOSHIMURA T *et al*. Leukocyte specificity and binding of human neutrophil attractant/activating protein 1. *J Immunol* 1990; **144**: 1323–1330.
  - 111 MATSUSHIMA K, LARSEN CG, DuBOIS GC, OPPENHEIM JJ. Purification and characterization of a novel chemotactic and activating factor produced by a human myelomonocytic cell line. *J Exp Med* 1989; **169**: 1485–1490.
  - 112 SCHALL TJ, BACON K, TOY KI, GOEDDEL DV. Selective attraction of monocytes and T lymphocytes of the memory phenotype by cytokine RANTES. *Nature* 1990; **34**: 769–771.
  - 113 BOYDEN S. The chemotactic effect of mixtures of antibody and antigen on polymorphonuclear leukocytes. *J Exp Med* 1962; **115**: 453–466.
  - 114 CUTLER JE. A simple *in vitro* method for the study of chemotaxis. *Proc Soc Exp Biol Med* 1974; **147**: 471–474.
  - 115 KELLER HU, WILKINSON PC, ABERCROMBIE M *et al*. A proposal for the definition of terms related to locomotion of leukocytes and other cells. *Clin Exp Immunol* 1977; **27**: 377–380.
  - 116 PEIPER SC, WANG ZX, NEOTE K *et al*. The Duffy antigen/receptor for chemokines (DARC) is expressed in endothelial cells of Duffy negative individuals who lack the erythrocyte receptor. *J Exp Med* 1995; **181**: 1311–1317.
  - 117 HORUK R. The interleukin-8-receptor family from chemokines to malaria. *Immunol Today* 1994; **15**: 169–174.
  - 118 BROADDUS VC, HEBERT CA, VITANGCOL RV *et al*. Interleukin-8 is a major neutrophil chemotactic factor in pleural liquid of patients with emphysema. *Am Rev Respir Dis* 1992; **146**: 825–830.
  - 119 KOCH AE, KUNKEL SL, BURROWS JC *et al*. Synovial tissue macrophages as a source of the chemotactic cytokine IL-8. *J Immunol* 1991; **147**: 2187–2195.
  - 120 STRIETER RM, KOCH AE, ANTONY VB *et al*. The immunopathology of chemotactic cytokines: the role of interleukin-8 and monocyte chemoattractant protein-1. *J Lab Clin Med* 1994; **123**: 183–197.
  - 121 ZEILLEMAKER AM, MUL FP, HOYNCK VAN PAPENDRECHT AA *et al*. Polarized secretion of interleukin-8 by human mesothelial cells: a role in neutrophil migration. *Immunology* 1995; **84**: 227–232.
  - 122 TAN P, HOMER-VANNIASINKAM S, GIMBRONE JR MA, LUSCINSKAS FW. Role of chemoattractants in neutrophil recruitment to sites of information. In: Paul LC, Issekutz TB, eds. *Adhesion Molecules in Health and Disease*. New York: Marcel Dekker, 1997: 257–287.
  - 123 LEY K, GAEHTGENS P. Endothelial, not hemodynamic differences are responsible for preferential leukocyte rolling in venules. *Circ Res* 1991; **69**: 1034–1041.
  - 124 PERRY MA, GRANGER DN. Role of CD11/CD18 in shear rate-dependent leukocyte-endothelial cell interactions in cat mesenteric venules. *J Clin Invest* 1991; **87**: 1798–1804.
  - 125 LAWRENCE MB, SMITH CW, ESKIN SG, McINTYRE LV. Effect of venous shear stress on CD18-mediated neutrophil adhesion to cultured endothelium. *Blood* 1990; **75**: 227–237.
  - 126 MOVAT HZ. The vascular phenomena of acute inflammation. In: Movat HZ, ed. *The Inflammatory Reaction*. Amsterdam: Elsevier, 1985: 43–75.
  - 127 SCHMID-SCHONBEIN GW, USAMI S, SKALAK R, CHIEN S. The interaction of leukocytes and erythrocytes in capillary and post-capillary vessels. *Microvasc Res* 1980; **19**: 45–70.
  - 128 NOBIS U, PRIES AR, COKELET GR, GAEHTGENS P. Radial distribution of white cells during blood flow in small tubes. *Microvasc Res* 1985; **29**: 295–304.
  - 129 TOZEREN A, LEY K. How do selectins mediate leukocyte rolling in venules? *Biophys J* 1992; **63**: 700–709.
  - 130 LICHTMAN AH, DING H, HENNAULT L *et al*. CD45RA<sup>+</sup>RO<sup>+</sup> (memory) but not CD45RA<sup>+</sup>RO<sup>-</sup> (naive) T cells roll efficiently on E- and P-selectin and VCAM-1 under flow. *J Immunol* 1997; **158**: 3640–3650.
  - 131 ATHERTON A, BORN GV. Quantitative investigations of the adhesiveness of circulating polymorphonuclear leucocytes to blood vessels. *J Physiol* 1972; **222**: 447–474.
  - 132 ATHERTON A, BORN GV. Relationship between the velocity of rolling granulocytes and that of the blood flow in venules. *J Physiol* 1973; **233**: 157–165.
  - 133 FINGER EB, PURI KD, ALON R *et al*. Adhesion through L-selectin requires a threshold hydrodynamic shear. *Nature* 1996; **379**: 266–269.
  - 134 GRANGER DN, KUBES P. The microcirculation and inflammation modulation of leukocyte-endothelial cell adhesion. *J Leuk Biol* 1994; **55**: 662–675.
  - 135 BARGATZE RF, KURK S, BUTCHER EC, JUTILA MA. Neutrophils roll on adherent neutrophils bound to cytokine-induced endothelial cells via L-selectin on the rolling cells. *J Exp Med* 1994; **180**: 1785–1792.
  - 136 ALON R, FUHLBRIGGE RC, FINGER EB, SPRINGER TA. Interactions through L-selectin between leukocytes and adherent leukocytes nucleate rolling adhesions on selectins and VCAM-1 in shear flow. *J Cell Biol* 1996; **135**: 849–865.
  - 137 ERLANDSEN SL, HASSLEN SR, NELSON RD. Detection and spatial distribution of the  $\beta_2$  integrin (Mac-1) and L-selectin (LECAM-1) adherence receptors on human neutrophils by high-resolution field emission SEM. *J Histochem Cytochem* 1993; **41**: 327–333.
  - 138 HUBER AR, KUNKEL SL, TODD III RF, WEISS SJ. Regulation of transendothelial neutrophil migration by endogenous interleukin-8. *Science* 1991; **254**: 99–102.
  - 139 TANAKA Y, ALBELDA SM, HORGAN KJ *et al*. CD31 expressed on distinctive T cell subsets is a preferential amplifier of  $\beta_1$  integrin-mediated adhesion. *J Exp Med* 1992; **176**: 245–253.
  - 140 GOODMAN RB, WOOD RG, MARTIN TR, HANSON-PAINTON O, KINASEWITZ GT. Cytokine-stimulated human mesothelial cells produce chemotactic activity for neutrophils including NAP-1/IL-8. *J Immunol* 1992; **148**: 457–465.
  - 141 ROLFE MW, KUNKEL SL, STANDIFORD TJ *et al*. Pulmonary fibroblast expression of interleukin-8: a model for alveolar macrophage-derived cytokine networking. *Am J Respir Cell Mol Biol* 1991; **5**: 493–501.
  - 142 TOPLEY N, BROWN Z, JORRES A *et al*. Human peritoneal mesothelial cells synthesize interleukin-8. Synergistic induction by interleukin-1-beta and tumor necrosis factor-alpha. *Am J Pathol* 1993; **142**: 1876–1886.
  - 143 JAFFE EA, GRULICH J, WEKSLER BB, HAMPEL G, WATANABE K. Correlation between thrombin-induced prostacyclin production and inositol triphosphate and cytosolic free calcium levels in cultured human endothelial cells. *J Biol Chem* 1987; **262**: 8557–8565.
  - 144 IVEY CL, WILLIAMS FM, COLLINS PD, JOSE PJ, WILLIAMS TJ. Neutrophil chemoattractants generated in two-phases during reperfusion of ischemic myocardium in the rabbit. *J Clin Invest* 1995; **95**: 2720–2728.
  - 145 McDONALD PP, POULIOT M, BERGEAT P, MCCOLL SR. Induction of chemokines of lipid mediator synthesis in granulocyte-macrophage colony-stimulating factor-treated human neutrophils. *J Immunol* 1993; **151**: 6399–6409.
  - 146 LOCATI M, ZHOU D, LUINI W, EVANGELISTA V, MANTOVANI A, SOZZANI S. Rapid induction of arachidonic acid release by monocyte chemoattractant protein-1 and related chemokines. Role of  $Ca^{2+}$  influx, synergism with platelet-activating factor and significance for chemotaxis. *J Biol Chem* 1994; **269**: 4746–4753.
  - 147 ROSS R. The pathogenesis of atherosclerosis: a perspective for the 1990s. *Nature* 1993; **362**: 801–809.
  - 148 FAGGIOTTO A, ROSS R, HARKER L. Studies of hypercholesterolaemia in the non-human primates. Changes that lead to fatty streak formation. *Arteriosclerosis* 1984; **4**: 323–340.
  - 149 LI H, CYBULSKY MI, GIMBRONE JR MA, LIBBY P. An atherogenic diet rapidly induces VCAM-1, a cytokine-regulated mononuclear leukocyte adhesion molecule, in rabbit aortic endothelium. *Arterioscler Thromb* 1993; **13**: 197–204.



- 150 STEMME S, HOLM J, HANSSON GK. T-lymphocytes in human atherosclerotic plaques are memory cells expressing CD45RO and the integrin VLA-1. *Arterioscler Thromb* 1992; **12**: 206–211.
- 151 DAVIES MJ, GORDON JL, GEARING AJ, PIGGOTT R, WOOLF N, KATZ D, KYRIAKOPOULOS A. The expression of the adhesion molecules ICAM-1, VCAM-1, PECAM and E-selectin in human atherosclerotic plaques. *J Pathol* 1993; **171**: 223–229.
- 152 JOHNSON-TIDEY RR, MCGREGOR JL, TAYLOR PR, POSTON RN. Increase in the adhesion molecule P-selectin in endothelium overlying atherosclerosis plaques. Co-expression with intercellular adhesion molecule-1. *Am J Pathol* 1994; **144**: 952–961.
- 153 WEYRICH AS, MCINTYRE TM, MCEVER RP, PRESCOTT SM, ZIMMERMAN GA. Monocyte tethering by P-selectin regulates monocyte chemotactic protein-1 and tumor necrosis factor- $\alpha$  secretion. *J Clin Invest* 1995; **95**: 2297–2303.
- 154 CUSHING SD, BERLINER JA, VALENTE AJ *et al*. Minimally modified low density lipoprotein induces monocyte chemoattractant protein-1 in human endothelial cells and smooth muscle cells. *Proc Natl Acad Sci USA* 1990; **87**: 5134–5138.
- 155 KU G, THOMAS CE, AKESON AL, JACKSON RL. Induction of interleukin-1 $\beta$  expression from human peripheral blood monocyte-derived macrophages by 9-hydroxyoctadecadienoic acid. *J Biol Chem* 1992; **267**: 14183–14188.
- 156 THOMAS CE, JACKSON RL, OHLEILER DE, KU G. Multiple lipid oxidation products in low density lipoproteins induce interleukin-1 $\beta$  release from human blood mononuclear cells. *J Lipid Res* 1994; **35**: 417–427.
- 157 LIBBY P, WARNER SJ, FRIEDMAN GB. Interleukin-1: A mitogen for human vascular smooth muscle cells that induces the release of growth-inhibitory prostanoids. *J Clin Invest* 1988; **81**: 487–498.
- 158 CLINTON SK, UNDERWOOD R, HAYES L *et al*. Macrophage colony-stimulating factor gene expression in vascular cells and in experimental and human atherosclerosis. *Am J Pathol* 1992; **140**: 301–316.
- 159 BROWNLEE M, CERAMI A, VLISSARA H. Advanced glycation end products in tissues and the biochemical basis of diabetic complications. *N Engl J Med* 1988; **318**: 1315–1321.
- 160 KIRSTEIN M, BRETT J, RADOFF S *et al*. Advanced protein glycosylation induced transendothelial human monocyte chemotaxis and secretion of platelet-derived growth factor: role in vascular disease of diabetes and aging. *Proc Natl Acad Sci USA* 1990; **87**: 9010–9014.
- 161 READ MA, NEISH AS, LUSCINSKAS FW *et al*. The proteasome pathway is required for cytokine-induced endothelial leukocyte adhesion molecule expression. *Immunity* 1995; **2**: 493–506.
- 162 PIERCE JW, SCHOENLEBER R, JESMOK G *et al*. Novel inhibitors of cytokine-induced I $\kappa$ B $\alpha$  phosphorylation and endothelial cell adhesion molecule expression show anti-inflammatory effects in vivo. *J Biol Chem* 1997; **272**: 21096–21103.
- 163 VEDDER NB, WINN RK, RICE CL *et al*. A monoclonal antibody to the adherence promoting leukocyte glycoprotein CD18 reduces organ injury and improves survival from hemorrhagic shock and resuscitation in rabbits. *J Clin Invest* 1988; **81**: 939–944.
- 164 KUBES P, JUTILA M, PAYNE D. Therapeutic potential of inhibiting leukocyte rolling in ischaemia/reperfusion. *J Clin Invest* 1995; **95**: 2510–2519.
- 165 SHARAR SR, WINN RK, MURRY CE, HARLAN JM, RICE CL. A CD18 monoclonal antibody increases the incidence and severity of subcutaneous abscess formation after high dose *S. aureus* injection in rabbits. *Surgery* 1991; **116**: 213–219.
- 166 MILESKE WJ, RAYMOND JE, WINN RK, HARLAN JM, RICE CL. Transient inhibition of neutrophil adherence with the anti-CD18 monoclonal antibody 60.3 does not increase mortality rates in abdominal sepsis. *Surgery* 1991; **109**: 497–501.
- 167 WALKER PM. Ischemia/reperfusion injury in skeletal muscle. *Ann Vasc Surg* 1991; **5**: 399–402.
- 158 WILLIAMS FM. Role of neutrophils in reperfusion injury. In: Hellewell PJ, Williams T, eds. *Immunopharmacology of Neutrophils*. London: Academic Press, 1994: 245–257.
- 169 WELBOURN CR, GOLDMAN G, PATERSON IS *et al*. Pathophysiology of ischemia reperfusion injury: central role of the neutrophil. *Br J Surg* 1991; **78**: 651–655.
- 170 WALDEN DL, MCCUTCHAN HJ, ENQUIST EJ *et al*. Neutrophils accumulate and contribute to skeletal muscle dysfunction after ischemia-reperfusion. *Am J Physiol* 1990; **259**: H1809–H1812.
- 171 SISLEY AC, DESAI T, HARIG JM, GEWERTZ BL. Neutrophil depletion attenuates human intestinal reperfusion injury. *J Surg Res* 1994; **57**: 192–196.
- 172 CRINNION JN, HOMER-VANNIASINKAM S, PARKIN SM, GOUGH MJ. Role of neutrophil-endothelial adhesion in skeletal muscle reperfusion injury. *Br J Surg* 1996; **83**: 251–254.
- 173 MEHTA J. Platelets and prostaglandins in coronary artery disease: rationale for use of platelet-suppressive drugs. *JAMA* 1983; **249**: 2818–2823.
- 174 CAPRIE STEERING COMMITTEE. A randomised, blinded, trial of clopidogrel versus aspirin in patients at risk of ischemic events (CAPRIE). *Lancet* 1996; **348**: 1329–1339.
- 175 COUKELL AJ, MARKHAM A. Clopidogrel. *Drugs* 1997; **54**: 745–750.
- 176 TOPOL EJ, CALIFF RM, WEISMAN HF *et al*. Randomised trial of coronary intervention with antibody against platelet IIb/IIIa integrin for reduction of clinical restenosis: results at six months. *Lancet* 1994; **343**: 881–886.
- 177 THE EPIC INVESTIGATORS. Use of a monoclonal antibody directed against the platelet glycoprotein IIb/IIIa receptor in high-risk coronary angioplasty. *N Engl J Med* 1994; **330**: 956–961.
- 178 THE EPILOG INVESTIGATORS. Platelet glycoprotein IIb/IIIa receptor blockade and low dose heparin during percutaneous coronary revascularisation. *N Engl J Med* 1997; **336**: 1689–1696.
- 179 FITZGERALD DJ, CATELLA F, ROY L, FITZGERALD GA. Marked platelet activation in vivo after intravenous streptokinase in patients with acute myocardial infarction. *Circulation* 1988; **77**: 142–150.
- 180 JANG I, GOLD HK, ZISKIND AA *et al*. Differential sensitivity of erythrocyte-rich and platelet-rich arterial thrombi to lysis with recombinant tissue-type plasminogen activator: a possible explanation for resistance to coronary thrombolysis. *Circulation* 1989; **79**: 920–928.
- 181 MILES LA, GINSBERG MH, WHITE JG, PLOW EF. Plasminogen interacts with human platelets through two distinct mechanisms. *J Clin Invest* 1986; **77**: 2001–2009.

Accepted 21 October 1998